

# References

References used as a source of information to construct the chart showing the 'Metabolic pathways of the diseased potato' are provided below.

## BIOSYNTHETIC PATHWAYS

[Ceramide](#)

[Cutin](#)

[Elicitors](#)

[Ethylene biosynthesis](#)

[Fatty acid degradation, lipoxygenase and oxylipin pathway](#)

[Glycoalkaloids](#)

[Halliwell-Asada pathway](#)

[Mevalonic acid pathway](#)

[Oxidative burst/free radicals](#)

[Pectin](#)

[Phenylpropanoid biosynthesis](#) (includes references on salicylic acid)

[Phosphoinositide cascade](#)

[Phosphorylation](#)

[Polyamine biosynthesis](#)

[PR proteins](#)

[Prenylation](#)

[Sesquiterpenes and phytoalexins](#)

[Shikimate pathway](#)

[Tropane alkaloids](#)

[Others](#)

## Ceramide

- Birch PRJ, Avrova AO, Duncan JM, Lyon GD, Toth RL. 1999. Isolation of potato genes which are induced during an early stage of the hypersensitive response to *Phytophthora infestans*. *Molecular Plant-Microbe Interactions* 12, 356-361. *Demonstrated that a number of genes, including serine palmitoyl-transferase, were up-regulated in response to infection by an incompatible isolate of P. infestans.*
- Brodersen P, Petersen M, Pike HM, Olszak B, Skov S, Odum N, Jorgensen LB, Brown RE, Mundy J. 2002. Knockout of Arabidopsis ACCELERATED-CELL-DEATH11 encoding a sphingosine transfer protein causes activation of programmed cell death and defense. *Genes Dev.* 16, 490-502.
- Da Silva D, Lachaud C, Cotelle V, Brière C, Grat S, Mazars C, Thuleau P. 2011. Nitric oxide production is not required for dihydrosphingosine-induced cell death in tobacco BY-2 cells. *May 1; 6(5).*
- Dickson RC. 1998. Sphingolipid functions in *Saccharomyces cerevisiae*: comparison to mammals. *Annual Review of Biochemistry* 67, 27-48
- Imai H, Ohnishi M, Kinoshita M, Kojima M, Ito S. 1995. Structure and distribution of cerebroside containing unsaturated hydroxy fatty acids in plant leaves. *Biosci.*

Biotech. Biochem. 59, 1309-1313

- Imai H, Ohnishi M, Hotsubo K, Kojima M, Ito S. 1997. Sphingoid base composition of cerebrosides from plant leaves. *Biosci. Biotech. Biochem.* 61, 351-353.
- Pena LA, Fuks Z, Kolesnick R. 1997. Stress-induced apoptosis and the sphingomyelin pathway. *Biochemical Pharmacology* 53, 615-621.

## **Cutin**

- Blee E, Schuber F. 1993. Biosynthesis of cutin monomers: involvement of a lipoxygenase/poroxygenase pathway. *The Plant Journal* 4, 113-123.
- Stapleton A, Beetham JK, Pinot F, Garbarino JE, Rockhold DR, Friedman M, Hammock BD, Belknap WR. 1994. Cloning and expression of soluble epoxidehydrolase from potato. *Plant Journal* 6, 251-258.

## **Elicitors**

- Chanda B, Xia Y, Mandal MK, Yu K, Sekine KT, Gao QM, Selote D, Hu Y, Stromberg A, Navarre D, Kachroo A, Kachroo P. 2011. Glycerol-3-phosphate is a critical mobile inducer of systemic immunity in plants. *Nat Genet* Mar 27.
- Cohen Y, Gisi U, Mosinger E. 1991. Systemic resistance of potato plants against *Phytophthora infestans* induced by unsaturated fatty acids. *Physiological and Molecular Plant Pathology* 38, 255-263.  
*Demonstrated that application of arachidonic, eicosapentaenoic, linoleic, linolenic and oleic acids to potato plants induced resistance to subsequent infection by P. infestans.*
- Cohen Y, Gisi U, Niderman T. 1993. Local and systemic protection against *Phytophthora infestans* induced in potato and tomato plants by jasmonic acid and jasmonic acid methyl ester. *Phytopathology* 83, 1054-1062.  
*Jasmonic acid (JA) and jasmonic acid methyl ester (JME) applied as foliar sprays to potato plants protected them from subsequent infection by P. infestans. Systemin did not induce protection. JA and JME application caused an increase in accumulation of proteinase inhibitors and other PR proteins in treated leaves.*
- Coquoz J-L, Buchala AJ, Meuwly P, Mettraux J-P. 1995. Arachidonic acid induces local but not systemic synthesis of salicylic acid and confers systemic resistance in potato plants to *Phytophthora infestans* and *Alternaria solani*. *Phytopathology* 85, 1219-1224.
- Du H, Klessig DF. 1997. Identification of a soluble, high-affinity salicylic acid-binding protein in tobacco. *Plant Physiology* 113, 1319-1327.  
*In contrast to earlier publications suggesting that catalase acts as a binding protein for salicylic acid this paper reports on the identification of a previously unreported*

*SA-binding protein.*

- Edwards R, Blount JW, Dixon RA. 1991. Glutathione and elicitation of the phytoalexin response in legume cell cultures. *Planta* 184, 403-409.  
*Both reduced glutathione and oxidised glutathione elicited phytoalexin biosynthesis via the phenylpropanoid pathway in cell-suspension cultures of Phaseolus vulgaris but had no effect in those of Medicago sativa. [There is no published evidence that glutathione, though a powerful elicitor in some plant species, induces a resistance-related response in potato].*
- Forrest RS, Lyon GD. 1990. Substrate degradation patterns of polygalacturonic acid lyase from *Erwinia carotovora* and *Bacillus polymyxa* and release of phytoalexin-eliciting oligosaccharides from potato cell walls. *Journal of Experimental Botany* 41, 481-488.  
*Phytoalexin elicitor-active oligogalacturonides released from pectin by the action of pectic enzymes produced by plant pathogens have now been reported in many systems. Only those pectin fragments of a particular size are effective phytoalexin elicitors. It is still not clear how many host responses are induced in plants by oligogalacturonides.*
- Kreft S, Ravnkar M, Mesko P, Pungercar J, Umek A, Kregar I, Strukelj B. 1997. Jasmonic acid inducible aspartic proteinase inhibitors from potato. *Phytochemistry* 44, 1001-1006.  
*Isolated four isoforms of aspartic proteinase inhibitors from potato and showed that mRNA was induced in potato shoots by 50 - 100  $\mu$ M jasmonic acid.*
- Laxalt AM, Cassia RO, Sanllorenti PM, Madrid EA, Andreu AB, Daleo GR, Conde RD, Lamattina L. 1996. Accumulation of cytosolic glyceraldehyde-3-phosphate dehydrogenase RNA under biological stress conditions and elicitor treatments in potato. *Plant Molecular Biology* 30, 961-972.  
*GAPDH gene transcripts increased in potato tubers treated with eicosapentaenoic acid and salicylic acid. (GAPDH is part of the glycolytic pathway in primary metabolism).*
- Maina G, Allen RD, Bhatia SK, Stelzig DA. 1984. Phenol metabolism, phytoalexins, and respiration in potato tuber tissue treated with fatty acid. *Plant Physiology* 76, 735-738.  
*Demonstrated that tuber discs treated with arachidonic acid became necrotic, accumulate sesquiterpene phytoalexins, and increases PAL activity and lignin accumulation.*
- Matsuda F, Miyagawa H, Ueno T. 2000. B-1,3-glucooligosaccharide induced activation of four enzymes responsible for N-p-coumaroyloctopamine biosynthesis in potato (*Solanum tuberosum* cv.) tuber tissue. *Zeitschrift fur Naturforschung C* 55, 373-382.
- Matton DP, Constabel P, Brisson N. 1990. Alcohol dehydrogenase gene expression in potato following elicitor and stress treatment. *Plant Molecular Biology* 14, 775-783.  
*mRNA of the ADH was induced in stems by arachidonic acid or by anaerobiosis.*

- Rickauer M, Brodschelm W, Bottin A, Veronesi C, Grimal H, Esquerre-Tugaye MT. 1997. The jasmonate pathway is involved differentially in the regulation of different defence responses in tobacco cells. *Planta* 202, 155-162.  
*In tobacco, expression of defence-related genes encoding proteinase inhibitor II, hydroxyproline-rich or glycine-rich glycoproteins, glucanase and chitinase can be induced by methyl jasmonate. Expression of the sesquiterpene cyclase (5-epi-aristolochene synthase) gene was not induced by methyl jasmonate. The synthesis of jasmonate via the LOX pathway is only part of a complex regulatory mechanism for the onset of many, but not all, defence reactions. There is no direct evidence whether these results can be extrapolated to potato.*
- Vidal S, de Leon IP, Denecke J, Palva ET. 1997. Salicylic acid and the plant pathogen *Erwinia carotovora* induce defense genes via antagonistic pathways. *The Plant Journal* 11, 115-123.  
*Demonstrated that salicylic acid (SA) is not the signaling molecule leading to the early response of tobacco to E. carotovora. Erwinia elicitors (not characterised) antagonize the SA-mediated induction of PR genes, and SA inhibited the induction of PR genes elicited by E. carotovora.*
- Yoshioka H, Sugie K, Park HJ, Maeda H, Tsuda N, Kawakita K, Doke N. 2001. Induction of plant gp91 phox homolog by fungal cell wall, arachidonic acid, and salicylic acid in potato. *Molecular Plant-Microbe Interactions* 14, 725-36.

### **Ethylene biosynthesis**

- Chao Q, Rothenberg M, Solano R, Roman G, Terzaghi W, Ecker JR. 1997. Activation of the ethylene gas response pathway in Arabidopsis by the nuclear protein ETHYLENE-INSENSITIVE3 and related proteins. *Cell* 89, 1133-1144.  
*Suggests a model of the ethylene signaling pathway in Arabidopsis including the existence of a MAPK cascade mediating signaling between CTR1 and EIN2 and the involvement of a family of DNA-binding proteins (ethylene-responsive element-binding proteins; EREBPs) mediating ethylene response gene activation.*
- Destefano-Beltran LJC, Van Caeneghem W, Gielen J, Richard L, Van Montagu M, Van Der Straeten D. 1995. Characterization of three members of the ACC synthase gene family in *Solanum tuberosum* L. *Molecular and General Genetics* 246, 496-508.  
*Showed the genes were modulated by wounding.*
- Edwards R. 1996. S-adenosyl-L-methionine metabolism in alfalfa cell cultures following treatment with fungal elicitors. *Phytochemistry* 43, 1163-1169. *Studied enzymes involved in maintaining an active pool of SAM for transmethylation reactions.*
- Guo H, Ecker JR 2003. Plant responses to ethylene gas are mediated by SCF(EBF1/EBF2)-dependent proteolysis of EIN3 transcription factor. *Cell* 115, 667-77.
- Kende H. 1993. Ethylene biosynthesis. *Annual Review of Plant Physiology and Plant Molecular Biology* 44, 283-307.

- Kieber JJ. 1997. The ethylene signal transduction pathway in *Arabidopsis*. *Journal of Experimental Botany* 48, 211-218.
- Liang W-S, Wen J-Q, Liang H-G. 1997. Stimulation of ethylene production in aged potato tuber slices by salicylic acid. *Phytochemistry* 44, 221-223.
- Liu Y, Su L-Y, Yang SF. 1985. Ethylene promotes the capability to malonylate 1-aminocyclopropane-1-carboxylic acid and D-amino acids in preclimacteric tomato fruits. *Plant Physiology* 77, 891-895.  
*Ethylene stimulated the ability to malonylate ACC. Malonylation is thought to inactivate foreign and potential toxic substances such as D-amino acids or some herbicides.*
- Malerba M, Bianchetti R. 1997. Stimulation of ACC oxidase activity "in vivo" by different effectors is differentially sensitive to inhibitors. *Journal of Plant Physiology* 150, 406-413.  
*Suggested that two differently regulated types of ethylene production are present in cultured sycamore cells.*
- Miyazaki JH, Yang SF. 1987. The methionine salvage pathway in relation to ethylene and polyamine biosynthesis. *Physiologia Plantarum* 69, 366-370. *Provides a summary diagram of the methionine cycle showing the biosynthesis of ethylene in relation to polyamine biosynthesis.*
- Oetiker JH, Olson DC, Shiu OY, Yang SF. 1997. Differential induction of seven 1-aminocyclopropane-1-carboxylate synthase genes by elicitor in suspension cultures of tomato (*Lycopersicon esculentum*). *Plant Molecular Biology* 34, 275-286.  
*ACC synthase is encoded by a multigene family. In tomato cell suspension cultures three ACC synthase genes were strongly induced by elicitor, three others were constitutively expressed, and one was present at all times at a very high level. The abundance of several ACC synthase mRNAs studied was not strictly correlated with the rate of elicitor-induced ethylene production.*
- Ouaked F, Rozhon W, Lecourieux D, Hirt H. 2003. A MAPK pathway mediates ethylene signaling in plants. *The EMBO Journal* 22, 1282-1288.

### **Fatty acid degradation, Lipoxygenase, and Oxylipin pathway**

- Alborn HT, Turlings TCJ, Jones TH, Stenhagen G, Loughrin JH, Tumlinson JH. 1997. An elicitor of plant volatiles from beet armyworm oral secretion. *Science* 276, 945-949.  
*Compares products of the lipoxygenase pathways in plants and animals.*

- Aliferis KA, Jabaji S. FT-ICR/MS and GC-EI/MS metabolomics networking unravels global potato sprout's responses to *Rhizoctonia solani* infection. PLoS One 7(8):e42576 Epub Aug 3  
*Identified azelaic acid in infected potato sprouts.*
- Blee E. 1995. Oxygenated fatty acids and plant defenses. INFORM 6, 852-861.  
*Summarises the lipoxygenase pathway and shows the hydroperoxide lyase pathway with 13-hydroperoxylinolenic acid as a substrate.*
- Blee E, Joyard J. 1996. Envelope membranes from spinach chloroplasts are a site of metabolism of fatty acid hydroperoxides. Plant Physiology 110,445-454.  
*Provides a summary of the major metabolic pathways leading from fatty acid hydroperoxides.*
- Bostock RM, Yamamoto H, Ricker KE, Choi D, Ward BL. 1991. Specific induction of 5-lipoxygenase activity in potato tuber by arachidonic acid. Phytopathology 81,1140 (abstract).  
*The arachidonic acid induced increase in 5-LOX enzyme activity precedes or parallels increases in mRNA abundance for PAL and HMG-CoA reductase. Non-elicitor fatty acids do not induce 5-LOX.*
- Bryant D, Cummins I, Dixon DP, Edwards R. 2006. Cloning and characterization of a theta class glutathione transferase from the potato pathogen *Phytophthora infestans*. Phytochemistry.  
*The P. infestans glutathione peroxidase converts 9(S)-HPOT to the non-fungitoxic 9(S)-HOT.*
- Butovich IA, Reddy CC. 2001. Enzyme-catalyzed and enzyme-triggered pathways in dioxygenation of 1-monolinoleoyl-*rac*-glycerol by potato tuber lipoxygenase. Biochemica et Biophysica Acta 1546, 379-398.
- Butovich IA, Luk'yanova SM, Reddy CC. 2000. Oxidation of linoleyl alcohol by potato tuber lipoxygenase: kinetics and positional, stereo, and geometrical (cis, trans) specificity of the reaction. Archives Biochem Biophys 378, 65-77.
- Chamulitrat W, Hughes MF, Eling TE, Mason RP. 1991. Superoxide and peroxy radical generation from the reduction of polyunsaturated fatty acid hydroperoxides by soybean lipoxygenase. Archives of Biochemistry and Biophysics 290, 153-159.  
*Provides a scheme to show the iron oxidation states of lipoxygenase during the formation and the decomposition of the fatty acid hydroperoxides.*
- Dammann C, Rojo E, Sanchez-Serrano JJ. 1997. Abscisic acid and jasmonic acid activate wound-inducible genes in potato through separate, organ-specific signal transduction pathways. The Plant Journal 11, 773-782.  
*Showed that jasmonic acid (JA) induces the expression of different sets of genes in roots and leaves of potato plants. When roots of intact plants were treated with JA, high levels of proteinase inhibitor II (pin2), cathepsin D inhibitor, leucine aminopeptidase and threonine deaminase mRNAs accumulated in the systemic leaves.*

*However, in treated roots, very low, if any expression of these genes could be detected. In contrast, a novel root-specific pin2 homologue accumulated in the JA-treated root tissue which could not be detected in leaves. Application of okadaic acid and staurosporine indicated that a protein phosphorylation step is involved in the regulation of this differential response.*

- Esterbauer H, Schaur RJ, Zollner H. 1991. Chemistry and biochemistry of 4-hydroxynonenal, malonaldehyde and related aldehydes. *Free Radical Biology and Medicine* 11, 81-128.  
*Reviews the biological activity of some lipid aldehydes including 4-hydroxynonenal, 4-hydroxyhexenal and malonaldehyde. These compounds have not yet been isolated from diseased potato but are likely to be produced during lipid peroxidation. These types of compounds are highly reactive and may act as second messengers which disseminate and augment initial free radical events.*
- Farmer EE, Ryan CA. 1992. Octadecanoid precursors of jasmonic acid activate the synthesis of wound-inducible proteinase inhibitors. *The Plant Cell* 4, 129-134.  
*Suggested that the octadecanoid intermediates may participate in a lipid-based signalling system that activates proteinase inhibitor synthesis in response to insect and pathogen attack.*
- Fauconnier M-L, Williams TD, Marlier M, Welti R. 2003. Potato tuber phospholipids contain colneleic acid in the 2-position. *FEBS Letters* 538, 155-158.
- Gardner HW. 1998. 9-hydroxy-traumatol, a new metabolite of the lipoxygenase pathway. *Lipids* 33, 745-749.
- Gidda SK, Miersch O, Schmidt J, Wasternack C, Varin L. 2003. Biochemical and molecular characterization of a hydroxy-jasmonate sulfotransferase from *A. thaliana*. *Journal of Biological Chemistry* March 10.  
*Not yet found in potato, but the authors suggest that hydroxylation and sulfonation reactions might be components of a pathway that inactivates excess jasmonate in plants.*
- Gobel C, Feussner I, Schmidt A, Scheel D, Sanchez-Serrano J, Hamberg M, Rosahl S. 2000. Oxylipin profiling reveals the preferential stimulation of the 9-lipoxygenase pathway in elicitor-treated potato cells. *Journal of Biological Chemistry* 276, 6267-6273.
- Grechkin AN, Kuramshin RA, Safonova EY, Yefremov YJ, Latypov SK, Ilyasov AV, Tarchevsky IA. 1991. Double hydroperoxidation of alpha-linolenic acid by potato tuber lipoxygenase. *Biochim Biophys Acta* 1081, 79-84.
- Gundlach H, Zenk MH, 1998. Biological activity and biosynthesis of pentacyclic oxylipins: the linoleic acid pathway. *Phytochemistry* 47, 527-537.
- Hamberg M. 1999. An epoxy alcohol synthase pathway in higher plants: biosynthesis of antifungal trihydroxy oxylipins in leaves of potato. *Lipids* 34, 1131-1142.
- Hamberg M, Sanz A, Castresana C. 1999. *B*-oxidation of fatty acids in higher plants. Identification of a pathogen-inducible oxygenase (PIOX) as an alpha-dioxygenase and

biosynthesis of 2-hydroperoxylinolenic acid. *The Journal of Biological Chemistry* 274, 24503-24513.

- Hamberg M. 2000. New cyclopentenone fatty acids formed from linoleic and linolenic acids in potato. *Lipids* 35, 353-363.
- Imbusch R, Mueller MJ. 2000. Analysis of oxidative stress and wound-inducible dinor isoprostanes F1 (phytoprostanes F1) in plants. *Plant Physiology* 124, 1293-1303.
- Itoh, A, Howe GA. 2001. Molecular cloning of a divinyl ether synthase. Identification as a CYP74 cytochrome P-450. *Journal of Biological Chemistry* 276, 3620-3627.
- Kirichenko A, Li L, Morandi MT, Holian A. 1996. 4-Hydroxy-2-nonenal-protein adducts and apoptosis in murine lung cells after acute ozone exposure. *Toxicology and Applied Pharmacology* 141, 416-424.  
*Demonstrated that several lipid peroxidation products including HNE (4-hydroxy-2-nonenal) are highly toxic to animal cells. They are likely to be produced in plants where lipid peroxidation occurs such as in infected tissues. Their effects on pathogen or plant cells in diseased tissues has not been determined.*
- Koda Y, Ward JL, Beale MH. 1995. Biological activity of methyl 7-methyl-jasmonates. *Phytochemistry* 38, 821-823.
- Koo AJ, Cooke TF, Howe GA. 2011. Cytochrome P450 CYP94B3 mediates catabolism and inactivation of the plant hormone jasmonyl-L-isoleucine. *PNAS USA* May 16.
- Martin M, Leon J, Dammann C, Albar JP, Griffiths G, Sanchez-Serrano JJ. 1999. Antisense-mediated depletion of potato leaf omega3 fatty acid desaturase lowers linolenic acid content and reduces gene activation in response to wounding. *European Journal of Biochemistry* 262, 283-90.
- Matsuura H, Ohkubo Y, Yoshihara T. 2001. Occurrence of 11-hydroxyjasmonic acid glucoside in leaflets of potato plants (*Solanum tuberosum* L.). *Biosci. Biotechnol. Biochem.* 65, 378-382.
- Matsuura H, Yoshihara T. 2003. Metabolism of deuterium-labeled jasmonic acid and OPC 8:0 in the potato plant (*Solanum tuberosum* L.) *Bioscience Biotechnology and Biochemistry* 67, 1903-1907.
- Parchmann S, Mueller MJ. 1998. Evidence for the formation of dinor isoprostanes E1 from alpha-linolenic acid in plants. *The Journal of Biological Chemistry* 273, 32650-32655.
- Pinot F, Skrabs M, Compagnon V, Salaun JP, Benveniste I, Schreiber L, Durst F. 2000. omega-Hydroxylation of epoxy- and hydroxy-fatty acids by CYP94A1: possible involvement in plant defence. *Biochemical Society Transactions* 28, 867-870.



- Royo J, Vancanneyt G, Perez AG, Sanz C, Stormann K, Rosahl S, Sanchez-Serrano JJ. 1996. Characterization of three potato lipoxygenases with distinct enzymatic activities and different organ-specific and wound-regulated expression patterns. *Journal of Biological Chemistry* 271, 21012-21019.
- Sembdner G, Parthier B. 1993. The biochemistry and the physiological and molecular actions of jasmonates. *Annual Review of Plant Physiology and Plant Molecular Biology* 44, 569-589.  
*Provides structures for a large number of naturally occurring jasmonates including esterified and amino acid conjugated products. Most of these have not yet been isolated from potato and have not been drawn on the chart.*
- Seo HS, Song JT, Cheong JJ, Lee YH, Lee YW, Hwang I, Lee JS, Choi YD. 2001. Jasmonic acid carboxyl methyltransferase: a key enzyme for jasmonate-regulated plant responses. *Proceedings of the National Academy of Science USA* 98, 4788-93.
- Spiteller G. 1996. Enzymic lipid peroxidation - a consequence of cell injury? *Free Radic. Biol. Med.* 21, 1003-9.  
*Summarises the steps involved in lipid peroxidation in injured cells.*
- Stumpe M, Kandzia R, Gobel C, Rosahl S, Feussner I. 2001. A pathogen-inducible divinyl ether synthase (CYP74D) from elicitor-treated potato suspension cells(1). *FEBS Lett* 507, 371-6.
- Vick BA, Zimmerman DC. 1984. Biosynthesis of jasmonic acid by several plant species. *Plant Physiology* 75, 458-461.  
*Summarises the biosynthesis of jasmonic acid from linolenic acid.*
- Wang C, Chin C-K, Ho C-T, Hwang C-F, Polashock JJ, Martin CE. 1996. Changes of fatty acids and fatty acid-derived flavor compounds by expressing the yeast delta-9 desaturase gene in tomato. *Journal of Agriculture and Food Chemistry* 44, 3399-3402.  
*Provides a summary diagram of hexanal production from linoleic acid through lipoxygenase/hydroperoxide lyase pathway and the autoxidation of palmitoleic acid.*
- Weber H, Chetelat A, Caldelari D, Farmer EE. 1999. Fatty acid synthesis in late blight-diseased potato leaves. *The Plant Cell* 11, 485-493.  
*Identified colneleic and colneleic acids in potato leaves infected with P. infestans.*
- Yoshihara T, Amanuma M, Tsutsumi T, Okumura Y, Matsuura H, Ichihara A. 1996. Metabolism and transport of [2-14C] jasmonic acid in the potato plant. *Plant Cell Physiology* 37, 586-590.  
*Concluded that jasmonic acid is metabolized to tuberonic acid glucoside*
- Ziegler J, Hamberg M, Miersch O, Parthier B. 1997. Purification and characterization of allene oxide cyclase from dry corn seeds. *Plant Physiology* 114, 565-573.  
*Allene oxide cyclase catalyzes the cyclization of 12,13(S)-epoxy-9(Z),11,15(Z)-octadecatrienoic acid to 12-oxo-10,15(Z)-phytodienoic acid. The generation of 12-oxo-PDA from 13-HPOT occurs via two enzymes; allene oxide synthase (AOS; also called hydroperoxide dehydratase) which is associated with membranes, and allene*

oxide cyclase (AOC) which is soluble.

## Glycoalkaloids

- Arnqvist L, Dutta PC, Jonsson L, Sitbon F. 2003. Reduction of cholesterol and glycoalkaloid levels in transgenic potato plants by overexpression of a type 1 sterol methyltransferase cDNA. *Plant Physiology* 131, 1792-1799.  
*Provides a schematic diagram of proposed sterol and glycoalkaloid biosynthesis pathways.*
- Bergenstrahle A, Tillberg E, Jonsson L. 1992. Regulation of glycoalkaloid accumulation in potato tuber discs. *Journal of Plant Physiology* 140, 269-275. *Studied the activity of several enzymes involved in glycoalkaloid biosynthesis after wounding. These enzymes included UDP glucose: solanidine glucosyltransferase, S-adenosyl-L-methionine:cycloartenol methyltransferase, and UDP-glucose:sterol glucosyltransferase.*
- Friedman M, McDonald GM. 1997. Potato glycoalkaloids: chemistry, analysis, safety, and plant physiology. *Critical Reviews in Plant Sciences* 16, 55-132.  
*An excellent summary of the glycoalkaloids present in potatoes and provides a possible synthetic pathway from cholesterol and cholesteranol to the aglycones.*
- Kaneko K, Tanaka MW, Mitsuhashi H. 1976. Origin of nitrogen in the biosynthesis of solanidine by *Veratrum grandiflorum*. *Phytochemistry* 15, 1391-1393.
- Laurila J, Laakso I, Valkonen JPT, Hiltunen, R. Pehu, E. 1996. Formation of parental-type and novel glycoalkaloids in somatic hybrids between *Solanum brevidens* and *S. tuberosum*. *Plant Science* 118, 145-155.  
*Quantified solanidine and solanthrene in S. tuberosum and tomatidine in S. brevidens and detected demissidine as a novel aglycone in hybrids. The authors hypothesised that the hydrogenase from S. brevidens produced demissidine from solanidine in the hybrids.*
- Moehs CP, Allen PV, Friedman M, Belknap WR. 1997. Cloning and expression of solanidine UDP-glucose glucosyltransferase from potato. *The Plant Journal* 11, 227-236.  
*Cloned a solanidine glucosyltransferase (SGT) and showed that mRNA increased transiently upon wounding.*
- Sawai S, Ohyama K, Yasumoto S, Seki H, Sakuma T, Yamamoto T, Takebayashi Y, Kojima M, Sakakibara H, Aoki T, Muranaka T, Saito K, Umemoto N. 2014. Sterol side chain reductase 2 is a key enzyme in the biosynthesis of cholesterol, the common precursor of toxic steroidal glycoalkaloids in potato. *Plant Cell* Sept 12.
- Silhavy D, Szentesi A, Banfalvi Z. 1996. *Solanum chacoense* lines with different alkaloid contents - a potential source of genes involved in leptine synthesis. *Acta Agronomica Hungarica* 44, 113-120.  
*Demonstrated the presence of hydroxy-solanidine and acetyl-solanidine in S.*

*chacoense*.

- Vazquez A, Gonzalez G, Ferreira F, Moyna P, Kenne L. 1997. Glycoalkaloids of *Solanum commersonii* Dun. ex Poir. *Euphytica* 95, 195-201.  
*Details glycoalkaloids and aglycones present in S. commersonii. These differ from those found in S. tuberosum.*

### **Halliwell-Asada pathway**

- Price AH, Taylor A, Ripley S J, Griffiths, A., Trewavas, A. J. 1994. Oxidative signals in tobacco increase cytosolic calcium. *The Plant Cell* 6, 1301-1310.  
*Summarises the Halliwell-Asada enzyme pathway, the main route for free radical formation in photosynthetic leaves.*

### **Mevalonic acid pathway**

- Chappell J. 1995. Biochemistry and molecular biology of the isoprenoid biosynthetic pathway in plants. *Annual Review of Plant Physiology and Plant Molecular Biology* 46, 521-547.  
*Reviewed many aspects of the mevalonic acid pathway. Described a hypothetical model suggesting that classes of isoprenoids are synthesised by discrete metabolic channels. For example, entire metabolic channels for sterol and sesquiterpene biosynthesis are localised in the endoplasmic reticulum. Synthesis of other isoprenoids, such as carotenoids and long chain prenols of ubiquinone, require the coordinated activity of cytoplasmic and organellar biosynthetic channels with ubiquinone synthesised in the mitochondria and carotenoids etc in the chloroplasts.*
- Choi D, Bostock RM, Avdiushko S, Hildebrand DF. 1994. Lipid-derived signals that discriminate wound- and pathogen-responsive isoprenoid pathways in plants: methyl jasmonate and the fungal elicitor arachidonic acid induce different 3-hydroxy-3-methylglutaryl-coenzyme A reductase genes and antimicrobial isoprenoids in *Solanum tuberosum* L. *Proceedings of the National Academy of Science USA* 91, 2329-2333.  
*Used gene-specific probes to study different HMGR genes (hmg1 and hmg2). Showed that methyl jasmonate can induce hmg1 expression and glycoalkaloid accumulation after wounding, and, importantly, reduce the abundance of hmg2 transcripts. Thus arachidonate and jasmonate-responsive pathways are distinct in relation to HMGR gene expression and isoprenoid product accumulation.*
- Choi D, Ward BL, Bostock RM. 1992. Differential induction and suppression of potato 3-hydroxy-3-methylglutaryl coenzyme A reductase genes in response to *Phytophthora infestans* and to its elicitor arachidonic acid. *The Plant Cell* 4, 1333-1344.  
*Isolated three classes of cDNAs encoding HMGR (hmg1, hmg2 and hmg3) from potato. Demonstrated a suppression of hmg1 and an enhancement of hmg2 and hmg3 transcripts following elicitor treatment or inoculation with an incompatible race of P. infestans.*

- Daudonnet S, Karst F, Tourte Y. 1997. Expression of the farnesyl diphosphate synthase gene of *Saccharomyces cerevisiae* in tobacco. *Molecular Breeding* 3, 137-145.  
*Transformed tobacco with FDP synthase gene from S. cerevisiae. Total FDP synthase-specific activity increased 12-fold in transformed plants and correlated with an increase of both sterol and carotenoid biosynthesis.*
- Learned RM, Connolly EL. 1997. Light modulates the spatial patterns of 3-hydroxy-3-methylglutaryl coenzyme A reductase gene expression in *Arabidopsis thaliana*. *The Plant Journal* 11, 499-511.  
*Showed that the hmg1 promoter is differentially responsive to light in different organs and promoter activation by light deprivation is confined primarily to immature leaves. In contrast, expression of the hmg1 gene in roots is confined to the elongation zone and is not responsive to illumination.*
- Ohnuma S, Hirooka K, Ohto C, Nishino T. 1997. Conversion from archael geranylgeranyl diphosphate synthase to farnesyl diphosphate synthase. *The Journal of Biological Chemistry*. 272, 5192-5198.  
*Investigated the molecular basis of substrate specificity for FPP synthase and GGPP synthase.*
- Scolnik PA, Bartley GE, 1996. A table of some cloned plant genes involved in isoprenoid biosynthesis. *Plant Molecular Biology Reporter* 14, 305-319.  
*Described the enzymes involved in the biosynthesis of isoprenoids in plants.*
- Weissbecker B, Van Loon JJA, Posthumus MA, Bouwmeester HJ, Dicke M. 2000. Identification of volatile potato sesquiterpenoids and their olfactory detection by the two-spotted stinkbug *Perillus bioculatus*. *Journal of Chemical Ecology* 26, 1433-1445.
- Weissenborn DL, Denbow CJ, Laine M, Lang SS, Yang Z, Yu X, Cramer CL. 1995. HMG-CoA reductase and terpenoid phytoalexins: Molecular specialization within a complex pathway. *Physiologia Plantarum* 93, 393-400.  
*Reviews evidence from a number of plants that HMGRs are encoded by a small gene family.*
- Yang Z, Park H, Lacy GH, Cramer CL. 1991. Differential activation of potato 3-hydroxy-3-methylglutaryl coenzyme A reductase genes by wounding and pathogen attack. *The Plant Cell* 3, 397-405.  
*Demonstrated that one isogene of the HMGR family is pathogen activated and is distinct from isogene(s) that are wound activated.*
- Yokota T. 1997. The structure, biosynthesis and function of brassinosteroids. *Trends in Plant Science* 2, 137-143.  
*Shows some of the steps involved in the biosynthesis of sterols in plants.*

## **Oxidative burst/free radicals**

- Adam AL, Bestwick CS, Barna B, Mansfield JW. 1995. Enzymes regulating the accumulation of active oxygen species during the hypersensitive reaction of bean to *Pseudomonas syringae* pv. *phaseolicola*. *Planta* 197, 240-249.  
*Found five main isoforms of superoxide dismutase (SOD) in Phaseolus vulgaris which increased after inoculation with incompatible bacteria. Also showed the existence of four peroxidases.*
- Arce DP, Tonon C, Zanetti ME, Godoy AV, Hirose S, Casalongue. 2006. The potato transcriptional co-activator StMBF1 is up-regulated in response to oxidative stress and interacts with the TATA-box binding protein. *J. Biochem Mol Biol* 39, 355-360.
- Deighton N, Johnston DJ, Glidewell SM, Lyon GD, Goodman BA. 1992. The involvement of oxygen-derived free radicals in the resistant response of potato tubers to *Erwinia carotovora*. *Protoplasma* 171, 167-169.  
*Demonstrated a rapid oxygen-requiring production of free radicals by potato tubers in response to infection by E. carotovora. They detected, by electron paramagnetic resonance (EPR) spectroscopy, a single radical adduct which had spectral characteristics of a lipid-derived radical.*
- Doke N, Miura Y, Sanchez LM, Kawakita K. 1994. Involvement of superoxide in signal transduction: responses to attack by pathogens, physical and chemical shocks, and UV irradiation. In, *Causes of photooxidative stress and amelioration of defense systems in plants*. (Eds C.H.Foyer and P.M.Mullineaux). CRC Press, Boca Raton, Ann Arbor, London, Tokyo. 177-197.  
*Summarises the relationship between the hypersensitive response of potato to P. infestans and superoxide production.*
- Kobayashi M, Ohura I, Kawakita K, Yokota N, Fujiwara M, Shimamoto K, Doke N, Yoshioka H. 2007. Calcium-dependent protein kinases regulate the production of reactive oxygen species by potato NADPH oxidase. *Plant Cell* 19, 1065-80.
- Lamb C, Dixon RA. 1997. The oxidative burst in plant disease resistance. *Annual Review of Plant Physiology and Plant Molecular Biology* 48, 251-275.  
*Reviews the literature on the oxidative burst in plant-pathogen interactions and suggests it is a central component of an integrated signalling system also involving salicylic acid and perturbations of Ca<sup>2+</sup> which underlies the expression of disease resistance mechanisms.*
- Langenkamper G, Manac'h N, Broin M, Cuine S, Becuwe N, Kuntz M, Rey P. 2001. Accumulation of plastid lipid-associated proteins (fibrillin/CDSP34) upon oxidative stress, ageing and biotic stress in Solanaceae and in response to drought in other species. *Journal of Experimental Botany* 52, 1545-1554.  
*Fibrillin/CDSP34 is induced in potato by Erwinia chrysanthemi.*
- Mata CG, Lamattina L, Cassia RO. 2001. Involvement of iron and ferritin in the potato-*Phytophthora infestans* interaction. *European Journal of Plant Pathology* 107, 557-562.
- Mehdy MC, Sharma YK, Sathasivan K, Bays NW. 1996. The role of activated oxygen species in plant disease resistance. *Physiologia Plantarum* 98, 365-374.

*Reviews the evidence demonstrating that the production of activated oxygen species is one of the earliest responses of plants to infection.*

- Pobezhimova TP, Voinikov VK. 2000. Biochemical and physiological aspects of ubiquinone function. *Membr Cell Biol* 13, 595-602.  
*Suggests that ubiquinol (the reduced form of ubiquinone) functions as an antioxidant in cells.*
- Rojas CM, Senthil-Kumar M, Wang K, Ryu CM, Kaundal A, Mysore KS. 2012. Glycolate oxidase modulates reactive oxygen species-mediated signal transduction during nonhost resistance in *Nicotiana benthamiana* and *Arabidopsis*. *Plant Cell* Jan 27.  
*Not yet demonstrated for potato but because of its occurrence in Nicotiana it is highly likely.*
- Yun B-W, Feechan A, Yin M, Saidi NBB, Le Bihan T, Yu M, Moore JW, Kang J-G, Kwon E, Spoel SH, Pallas JA, Loake GJ. 2011. S-nitrosylation of NADPH oxidase regulates cell death in plant immunity. *Nature* 478, 264-8.

## **Pectin**

- Bourlard T, Schaumann-Gaudinet A, Bruyant-Vannier M-P, Morvan C. 1997. Various pectin methyltransferase activities with affinity for low and highly methylated pectins. *Plant Cell Physiology* 38, 259-267.
- Ishii T. 1997. O-acetylated oligosaccharides from pectins of potato tuber cell walls. *Plant Physiology* 113, 1265-1272.  
*Describes the detection of some acetylated galacturonides from potato cell walls.*
- McMillan GP, Hedley D, Fyffe L, Perombelon MCM. 1993. Potato resistance to soft rot erwinias is related to cell wall pectin esterification. *Physiological and Molecular Plant Pathology* 42, 279-289.  
*Using potato clones derived from *S. brevidens* showed a correlation between pectin esterification and resistance to stem rotting by *E. carotovora*: pectin from the more resistant clones was more highly methylated. More recent data (unpublished) suggests that there is not a simple relationship between resistance and pectin methylation.*

## **Phenylpropanoid biosynthesis**

- Belles JM, Garro R, Fayos J, Navarro P, Primo J, Conejero V. 1999. Gentisic acid as a pathogen-inducible signal, additional to salicylic acid for activation of plant defenses in tomato. *Molecular Plant-Microbe Interactions* 12, 227-235.  
*Detected gentisic acid in virus infected tomato.*
- Bernards MA, Lewis NG. 1992. Alkyl ferulates in wound healing potato tubers. *Phytochemistry* 31, 3409-3412.

*Showed that alkyl ferulate esters begin to accumulate 3 - 7 days after wounding potato tubers; these are associated with the suberin of wound periderm formation.*

- Bernards MA, Lopez ML, Zajicek J, Lewis NG. 1995. Hydroxycinnamic acid-derived polymers constitute the polyaromatic domain of suberin. *The Journal of Biological Chemistry* 270, 7382-7386.  
*Studied the phenolic domain of suberin and suggested that it was almost exclusively comprised of a covalently-linked hydroxycinnamic acid-derived polymeric matrix.*
- Brady JD, Fry SC. 1997. Formation of di-isodityrosine and loss of isodityrosine in the cell walls of tomato cell-suspension cultures treated with fungal elicitors or H<sub>2</sub>O<sub>2</sub>. *Plant Physiology* 115, 87-92  
*Treatment of tomato cells with resistance elicitors or H<sub>2</sub>O<sub>2</sub> led to increases in Tyr-based oxidative coupling in the plant cell wall with the formation of di-isodityrosine. (Similar reactions probably occur in potato).*
- Brady JD, Sadler IH, Fry SC. 1997. Pulcherosine, an oxidatively coupled trimer of tyrosine in plant cell walls: its role in cross-link formation. *Phytochemistry* 47, 349-353.  
*Isolated pulcherosine, a tyrosine-based trimer composed of isodityrosine and tyrosine oxidatively coupled via a biphenyl link, from tomato cell walls. Pulcherosine forms inter-polypeptide cross-links and/or wide intra-polypeptide loops.*
- Chong J, Pierrel M-A, Atanassova R, Werck-Reichhart D, Fritig B, Saindrenan P. 2001. Free and conjugated benzoic acid in tobacco plants and cell cultures. Induced accumulation upon elicitation of defense responses and role as salicylic acid precursors. *Plant Physiology* 2001. 125, 318-328.  
*The pathway proposed is for tobacco; it may be similar in potato though not yet proven.*
- Clarke DD. 1982. The accumulation of cinnamic acid amides in the cell walls of potato tissue as an early response to fungal attack. In, *Active Defense Mechanisms in Plants*. (ed, R. K. S. Wood). Plenum Press, New York, London.321-322.  
*Described the presence of amides of cinnamic acid and p-coumaric acid with tyramine or octopamine in potato tubers infected with P. infestans*
- Delaney TP. 1997. Genetic dissection of acquired resistance to disease. *Plant Physiology* 113, 5-12.  
*Salicylic acid is converted to catechol by salicylate hydroxylase which is encoded by the nahG gene from Pseudomonas putida. This pathway therefore only occurs in genetically transformed plants.*
- Dixon RA, Paiva NL. 1995. Stress-induced phenylpropanoid metabolism. *The Plant Cell* 7, 1085-1097.  
*Outlined the biosynthetic relationships among many classes of phenylpropanoids. Many phenolics accumulate in the vacuole as glycosides or other conjugates.*
- Douglas CJ. 1996. Phenylpropanoid metabolism and lignin biosynthesis: from weeds to trees. *Trends in Plant Science* 1, 171-178.  
*A summary of the biosynthesis of phenylpropanoids leading to the production of*

*lignin in plants.*

- El Hadrami A, Adam LR, Daayf F. 2011. Biocontrol treatments confer protection against *Verticillium dahlia* infection of potato by inducing anti-microbial metabolites. *Molecular Plant-Microbe Interactions* 24; 328-35.  
*Described the induction of rutin in potato and the detection of 2-PCPGCA.*
- Friedman M. 1997. Chemistry, biochemistry, and dietary role of potato polyphenols. A review. *Journal of Agriculture and Food Chemistry* 45, 1523-1540.  
*Describes analytical and compositional aspects of phenolic compounds in potatoes; their biosynthesis, molecular genetics, and role in resistance to plant pathogens. Proposes a biosynthetic route to chlorogenic acid from cinnamic acid via cinnamoyl-D-glucose to p-coumaroyl-D-glucose and caffeoyl-D-glucose. Proposes a scheme to explain inhibition of enzymatic browning by cysteine and ascorbic acid.*
- Henriquez MA, Adam LR, Daayf F. 2012. Alteration of secondary metabolites' profiles in potato leaves in response to weakly and highly aggressive isolates of *Phytophthora infestans*. *Plant Physiol Biochem* May 3.
- Jarvis AP, Schaaf O, Oldham NJ. 2000. 3-Hydroxy-3-phenylpropanoic acid is an intermediate in the biosynthesis of benzoic acid and salicylic acid but benzaldehyde is not. *Planta* 212, 119-126.
- Joos H-J, Hahlbrock K. 1992. Phenylalanine ammonia-lyase in potato (*Solanum tuberosum* L.). *European Journal of Biochemistry* 204, 621-629.  
*Suggested that potato contains approximately 40 - 50 PAL genes/haploid genome. cDNA heterogeneity suggests that possibly more than 10 of these genes are potentially active. Mature leaves contained particularly low levels of PAL mRNA but infection by *Phytophthora infestans* caused a large transient increase. Describes two gene families, PAL1 and PAL2.*
- Keller H, Hohlfeld H, Wray V, Hahlbrock K, Scheel D, Strack D. 1996. Changes in the accumulation of soluble and cell wall-bound phenolics in elicitor-treated cell suspension cultures and fungus-infected leaves of *Solanum tuberosum*. *Phytochemistry* 42, 389-396.  
*Used HPLC to quantify pathogen- and elicitor-induced changes of potato phenolics. The largest changes were shown by soluble putrescine amides and cell wall-bound 4-hydroxybenzaldehyde and tyramine amides. Large amounts of coumaroyltyramine were secreted into the culture medium.*
- King RR, Calhoun LA. 2005. Characterization of cross-linked hydroxycinnamic acid amides isolated from potato common scab lesions. *Phytochemistry* 66, 2468-2473.
- Kröner A, Marnet N, Andrivon D, Val F. 2012. Nicotiflorin, rutin and chlorogenic acid: phenylpropanoids involved differently in quantitative resistance of potato tubers to biotrophic and necrotrophic pathogens. *Plant Physiol Biochem* 15, 57C: 23-31.
- Lacombe E, Hawkins S, Van Doorselaere J, Piquemal J, Goffner D, Poeydomenge O, Boudet A-M, Grima-Pettenati J. 1997. Cinnamoyl CoA reductase, the first committed enzyme of the lignin biosynthetic pathway: cloning, expression and



phylogenetic relationships. *The Plant Journal* 11, 429-441.

*Cinnamoyl CoA:NADP oxidoreductase (CCR) catalyzes the conversion of cinnamoyl CoA esters to their corresponding cinnamaldehydes.*

- Lee H-i, Raskin, I. 1998. Glucosylation of salicylic acid in *Nicotiana tabacum* cv. Xanthi-nc. *Phytopathology* 88, 692-697.  
*Describes the activity of salicylic acid 2-O-glucosyltransferase, salicylic acid carboxyl glucosyltransferase and salicylic acid methyltransferase in infected tobacco.*
- Lyon GD, McGill FM. 1988. Inhibition of growth of *Erwinia carotovora* in vitro by phenolics. *Potato Research* 31, 461-467.  
*Showed that the phenolic acids caffeic, cinnamic, ferulic, salicylic, sinapic, and vanillic, together with scopoletin and coniferyl alcohol inhibited growth of E. carotovora in vitro. Chlorogenic acid did not inhibit growth of E. carotovora in buffered nutrient broth.*
- Malmberg A, Theander O. 1984. Free and conjugated phenolic acids and aldehydes in potato tubers. *Swedish Journal of Agricultural Research* 14, 119-125.  
*Used GC analysis to detect a wide range of free and conjugated phenolic acids and aldehydes in healthy potato tubers.*
- Manosalva PM, Park SW, Forouhar F, Tong L, Fry WE, Klessig DF. 2010. Methyl esterase 1 (StMES1) is required for systemic acquired resistance in potato. *Molecular Plant-Microbe Interactions* 23, 1151-63.
- Mayer AM, Harel E. 1979. Polyphenol oxidases in plants. *Phytochemistry* 18, 193-215.  
*Critically reviews catechol oxidases and laccases (both classified as polyphenol oxidases) in plants; their purification, subcellular localisation and properties.*
- Muhlenbeck U, Barz W. 1997. Cytochrome-P450-dependent formation of alpha-hydroxyacetovanillone from acetovanillone in *Solanum khasianum*. *Phytochemistry* 44, 865-867.  
*Studied the P450-dependent monooxygenase catalyzing the hydroxylation of acetovanillone.*
- Negrel J, Lotfy S, Javelle F. 1995. Modulation of the activity of two hydroxycinnamoyl transferases in wound-healing potato tuber discs in response to pectinase or abscisic acid. *Journal of Plant Physiology* 146, 318-322.  
*Tyramine hydroxycinnamoyl transferase (THT) activity was strongly stimulated in potato tubers by pectinase and, to a lesser extent, by ABA. In contrast - hydroxypalmitic acid hydroxycinnamoyl transferase (HHT), which is associated with suberin biosynthesis, was induced by ABA but not by pectinase. Neither arachidonic nor jasmonic acids affected induction of THT or HHT activities.*
- Negrel J, Pollet B, Lapierre C. 1996. Ether-linked ferulic acid amides in natural and wound periderms of potato tuber. *Phytochemistry* 43, 1195-1199.  
*Used thioacidolysis and desulphuration followed by GC-MS to demonstrate the occurrence of feruloyltyramine and /or feruloyloctopamine in natural and wound periderm of potato tubers.*

- Pallas JA, Paiva NL, Lamb C, Dixon RA. 1996. Tobacco plants epigenetically suppressed in phenylalanine ammonia-lyase expression do not develop systemic acquired resistance in response to infection by tobacco mosaic virus. *The Plant Journal* 10, 281-293.

*PR proteins were not induced in systemic leaves of PAL-suppressed plants. De novo synthesis of salicylic acid and/or synthesis of other phenylpropanoids is required for the expression of resistance to TMV in systemic leaves.*
- Razal RA, Ellis S, Singh S, Lewis NG, Towers GHN. 1996. Nitrogen recycling in phenylpropanoid metabolism. *Phytochemistry* 41, 31-35.

*Proposes a scheme for nitrogen recycling in phenylpropanoid metabolism. Ammonium ions released by PAL during the conversion of phenylalanine to cinnamic acid is recycled in the biosynthesis of arogenic acid from prephenic acid.*
- Rosler J, Krekel F, Amrhein N, Schmid J. 1997. Maize phenylalanine ammonia-lyase has tyrosine ammonia-lyase activity. *Plant Physiology* 113, 175-179.

*Expression of a PAL-specific cDNA from maize in E. coli proved that phenylalanine ammonia-lyase (PAL) and tyrosine ammonia-lyase (TAL) activities reside in one polypeptide (unlike dicots) and imply that maize can produce p-coumaric acid from both phenylalanine and tyrosine. Potato does not possess TAL.*
- Ryals JA, Neuenschwander UH, Willits MG, Molina A, Steiner H-Y, Hunt MD. 1996. Systemic acquired resistance. *The Plant Cell* 8, 1809-1819.

*Proposed two alternative routes, involving either -oxidation or a non-oxidative pathway, by which cinnamic acid could be converted into benzoic acid in plants. These routes have not been confirmed in potato.*
- Serra O, Figueras M, Franke R, Prat S, Molinas M. 2010. Unraveling ferulate role in suberin and periderm biology by reverse genetics. *Plant Signal Behav* 5; 953-958.
- Schmidt A, Grimm R, Schmidt J, Scheel D, Strack D, Rosahl S. 1999. Cloning and expression of a potato cDNA encoding hydroxycinnamoyl-CoA:tyramine N-(hydroxycinnamoyl)transferase. *The Journal of Biological Chemistry* 274, 4273-4280.

*Provides structures of the hydroxycinnamic acid amides accumulating in cell walls and medium of suspension cultured cells.*
- Schoch G, Goepfert S, Morant M, Hehn A, Meyer D, Ullmann P, Werck-Reichhart D. 2001. CYP98A3 from *Arabidopsis thaliana* is a 3'-hydroxylase of phenolic esters, a missing link in the phenylpropanoid pathway. *The Journal of Biological Chemistry* 276, 36566-36574.

*This enzyme has not yet been confirmed in potato.*
- Shulaev V, Silverman P, Raskin I. 1997. Airborne signalling by methyl salicylate in plant pathogen resistance. *Nature* 385, 718-721.

*Showed that methyl salicylate, produced in many plants, is a major volatile produced by tobacco plants inoculated with tobacco mosaic virus.*

- Stevens LH, Davelaar E, Kolb RM, Pennings EJM, Smit NPM. 1998. Tyrosine and cysteine are substrates for blackspot synthesis in potato. *Phytochemistry* 49, 703-707. *Describes the biosynthetic route for eumelanin and phaeomelanin*
- Szopa J, Wilczynski G, Fiehn O, Wenzel A, Willmitzer L. 2001. Identification and quantification of catecholamines in potato plants (*Solanum tuberosum*) by GC-MS. *Phytochemistry* 58, 315-320. *Identified dopamine, norepinephrine and normetanephrine.*
- Thipyapong P, Hunt MD, Steffens JC. 1995. Systemic wound induction of potato (*Solanum tuberosum*) polyphenol oxidase. *Phytochemistry* 40, 673-676. *Potato PPO can be systemically induced by wounding but only in tissues which normally express steady-state PPO mRNA.*
- Thygesen PW, Dry IB, Robinson SP. 1995. Polyphenol oxidase in potato. A multigene family that exhibits differential expression patterns. *Plant Physiology* 109, 525-531. *PPO catalyzes the conversion of monophenols to o-diphenols and o-dihydroxyphenols to o-quinones. The quinone products can then polymerize and react with amino groups of proteins causing black or brown pigments. PPO is present as a small multigene family in potato and each gene has a specific temporal and spacial pattern of expression.*
- van Gelder CWG, Flurkey WH, Wichers HJ. 1997. Sequence and structural features of plant and fungal tyrosinases. *Phytochemistry* 45, 1309-1323. *Provides a reaction scheme for tyrosinase showing both the cresolase and catecholase activity.*
- van Verk MC, Bol, JF, Linthorst HJ. 2011. WRKY transcription factors involved in activation of SA biosynthesis genes. *BMC Plant Biol* 11, 89.
- Wildermuth MC, Dewdney J, Wu G, Ausubel FM. 2001. Isochorismate synthase is required to synthesize salicylic acid for plant defense. *Nature* 414, 562-565. *Evidence is derived from experiments on Arabidopsis. There is no comparable experimental data yet for potato.*
- Zhang X-H, Chiang VL. 1997. Molecular cloning of 4-coumarate:coenzyme A ligase in loblolly pine and the roles of this enzyme in the biosynthesis of lignin in compression wood. *Plant Physiology* 113, 65-74. *Summarised the lignin biosynthetic pathway in gymnosperms.*

### **Phosphoinositide cascade**

- Lemtiri-Chlieh F, MacRobbie EAC, Brearley CA. 2000. Inositol hexakisphosphate is a physiological signal regulating the K<sup>+</sup>-inward rectifying conductance in guard cells. *Proc. Natl. Acad. Sci. USA* 97, 8687-8692.
- Fruman DA, Meyers RE, Cantley LC. 1998. Phosphoinositide kinases. *Annual Review of Biochemistry* 67, 481-507.

- Liscovitch M, Cantley LC. 1994. Lipid second messengers. *Cell* 77, 329-334.  
*Reviews information on signalling pathways of lipid second messengers. Most of the information concerns animal cells.*
- Lupu VD, Kaznatcheyeva E, Krishna UM, Falck JR, Bezprozvanny I. 1998. Functional coupling of phosphatidylinositol 4,5-bisphosphate to inositol 1,4,5-trisphosphate receptor. *The Journal of Biological Chemistry* 273, 14067-14070.
- Munnik T. 2001. Phosphatidic acid: an emerging plant lipid second messenger. *Trends in Plant Science* 6, 227-233.
- Munnik T, de Vrije T, Irvine RF, Musgrave A. 1996. Identification of diacylglycerol pyrophosphate as a novel metabolic product of phosphatidic acid during G-protein activation in plants. *The Journal of Biological Chemistry* 271, 15708-15715.
- Nakano T, Joh T, Narita K, Hayakawa T. 2000. The pathway of dephosphorylation of myo-inositol hexakisphosphate by phytases from wheat bran of *Triticum aestivum* L. cv. Nourin #61. *BioSci. Biotechnol. Biochem.* 64, 995-1003.
- Poovaiah BW, Reddy ASN, McFadden JJ. 1987. Calcium messenger system: Role of protein phosphorylation and inositol bisphospholipids. *Physiologia Plantarum* 69, 569-573.  
*Reviews information on the role of inositol phospholipids in signal transduction in plants.*
- Raboy V. 2003. myo-Inositol-1,2,3,4,5,6-hexakisphosphate. *Phytochemistry* 64, 1033-1043.
- Riedel B, Morr M, Wu WI, Carman GM, Wissing JB. 1997. Metabolism of diacylglycerol pyrophosphate by suspension cultured *Catharanthus roseus* cells - identification and characterization of diacylglycerol pyrophosphatase in plants. *Plant Science* 128, 1-10.  
*Identification of DGPP phosphatase in a wide range of organisms.*
- Saiardi A, Caffrey JJ, Snyder SH, Shears SB. 2000. The inositol hexakisphosphate kinase family. Catalytic flexibility and function in yeast vacuole biogenesis. *The Journal of Biological Chemistry* 275, 24686-24692.
- Wissing JB, Behrbohm H. 1993. Phosphatidate kinase, a novel enzyme in phospholipid metabolism. Purification, subcellular localization, and occurrence in the plant kingdom. *Plant Physiology* 102, 1243-1249.

## **Phosphorylation**

- Arino J, Perez-Callejon E, Cunillera N, Camps M, Posas F, Ferrer A. 1993. Protein phosphatases in higher plants: multiplicity of type 2A phosphatases in *Arabidopsis thaliana*. *Plant Molecular Biology* 21, 475-485.  
*The authors suggest that a multiplicity of type 2A phosphatases might be differentially expressed in higher plants.*

- Baizabal-Aguirre VM, de la Vara LEG. 1997. Purification and characterization of a calcium-regulated protein kinase from beet root (*Beta vulgaris*) plasma membranes. *Physiologia Plantarum* 99, 135-143.  
*Purified a calcium-regulated protein kinase (CRPK) associated with the plasma membrane of beet root cells. Calcium regulated kinases are likely to be present in all plants including potato.*
- Bianchini GM, Stermer BA, Paiva NL. 1996. Induction of early mevalonate pathway enzymes and biosynthesis of end products in potato (*Solanum tuberosum*) tubers by wounding and elicitation. *Phytochemistry* 42, 1563-1571.  
*HMGR levels increased 30-fold following arachidonic acid treatment and 15-fold following wounding, but mevalonate kinase and mevalonate phosphate kinase only increased two- to four-fold following these treatments. While HMGR levels are extremely low in untreated tissues, the activities of the two kinases are relatively high suggesting to the authors that they do not serve as control points for the synthesis of terpenoids.*
- Bolwell GP. 1992. A role for phosphorylation in the down-regulation of phenylalanine ammonia-lyase in suspension-cultured cells of French bean. *Phytochemistry* 31, 4081-4086.  
*Provided evidence that PAL activity in Phaseolus vulgaris is (partially) controlled by phosphorylation. [There is no comparable evidence concerning phosphorylation of PAL in potato but it seems likely that there will be some similarities].*
- Carcamo JM, Pedraza A, Borquez-Ojeda O, Zhang B, Sanchez R, Golde DW. 2004. Vitamin C is a kinase inhibitor: dehydroascorbic acid inhibits I $\kappa$ B kinase beta. *Molecular and Cellular Biology* 24, 6645-6652.  
*Work done on HeLa cells. No published evidence yet for plant cells*
- Conrath U, Silva H, Klessig DF. 1997. Protein dephosphorylation mediates salicylic acid-induced expression of PR-1 genes in tobacco. *The Plant Journal* 11,747-757.  
*Using protein kinase and phosphatase inhibitors they showed that PR1 gene induction in tobacco can be mediated by dephosphorylation of serine/threonine residue(s) of two or more unidentified phosphoproteins. Protein phosphatase inhibitors okadaic acid and calyculin A blocked SA-mediated induction of PR-1 genes implying the involvement of a phosphoprotein downstream of SA. The protein kinase inhibitors K-252a and staurosporine induced PR-1 gene expression. K-252a also stimulated production of SA and its glucoside, suggesting that another phosphoprotein acts upstream of SA. These results suggest that there are two (or more) phosphoproteins which function in the same signal transduction pathway leading to PR-1 gene induction.*
- Felix G, Grosskopf DG, Regenass M, Boller T. 1991. Rapid changes of protein phosphorylation are involved in transduction of the elicitor signal in plant cells. *Proceedings of the National Academy of Science USA* 88, 8831-8834.  
*Showed that phosphorylation of protein substrates by K-252a-sensitive protein kinases is essential for transduction of elicitor signals in tomato cells and continuous phosphorylation of these proteins is required to maintain the elicited response.*

- Hart GW, Kreppel LK, Comer FI, Arnold CS, Snow DM, Ye Z, Cheng X, Della Manna D, Caine DS, Earles BJ, Akimoto Y, Cole RN, Hayes BK. 1996. O-GlcNAcylation of key nuclear and cytoskeletal proteins: reciprocity with O-phosphorylation and putative roles in protein multimerization. *Glycobiology* 6, 711-716.  
*O-GlcNAcylated proteins are ubiquitous in all eukaryotes examined. The authors suggest a possible regulatory modification of some proteins involving a reciprocal relationship between phosphorylation of a hydroxyl moiety and its O-GlcNAcylation. If the system is truly widespread amongst eukaryotes then it is likely to be present in potato although there is no direct evidence.*
- Hassan A, Okuta T, Kato M, Hatsugai N, Sano Y, Ishimori T, Okazaki K, Doullah MA, Shah MM. 2012. Alternaric acid stimulates phosphorylation of His-tagged RiCDPK2, a calcium-dependent protein kinase in potato plants. *Genetics and Molecular Research* May 10.
- Hirt H. 1997. Multiple roles of MAP kinases in plant signal transduction. *Trends in Plant Science* 2, 11-15.  
*Reviews the literature on MAP kinases involved in signal transduction caused by wounding, pathogens and abiotic stresses as well as the hormones abscisic acid, auxin and ethylene. Thus, pathways involving MAP kinases are common in signal transduction in all eukaryotes. MAP (mitogen-activated protein) kinases are serine/threonine protein kinases originally described in animals.*
- Jonak C, Heberle-Bors E, Hirt H. 1994. MAP kinases: universal multi-purpose signaling tools. *Plant Molecular Biology* 24, 407-416.  
*Demonstrated the presence of MAP kinase, MAP kinase kinase and MAP kinase kinase kinase in plants and suggested that they appear to be involved in a number of physiological responses including ethylene signaling and auxin-induced cell proliferation.*
- Katou S, Senda K, Yoshioka H, Doke N, Kawakita K. 1999. A 51 kDa protein kinase of potato activated with hyphal wall components from *Phytophthora infestans*. *Plant Cell Physiol* 40, 825-831.
- Kim JH, Kim WT, Kang BG, Yang SF. 1997. Induction of 1-aminocyclopropane-1-carboxylate oxidase mRNA by ethylene in mung bean hypocotyls: involvement of both protein phosphorylation and dephosphorylation in ethylene signalling. *The Plant Journal* 11, 399-405.  
*The protein kinase inhibitor staurosporine prevented ethylene-induced ACC oxidase gene expression whereas it recovered the ethylene-suppressed transcript level of ACC synthase suggesting that protein phosphorylation plays a role in the induction of ACC oxidase and suppression of ACC synthase by ethylene in mung bean hypocotyls. Biosynthesis of ethylene appears to occur via the same pathway in all higher plants and control through phosphorylation is therefore also likely to occur in potato.*
- MacKintosh RW, MacKintosh C. 1993. Regulation of plant metabolism by reversible protein (serine/threonine) phosphorylation. In, *Post-translational modifications in plants*, ed. N.H. Battey, H.G. Dickinson. Society for Experimental Biology Seminar Series 53. Cambridge University Press. 197-212.

*Describes the protein kinase cascade controlling HMGR.*

- MacKintosh RW, Davies SP, Clarke PR, Weekes J, Gillespie JG, Gibb BJ, Hardie DG. 1992. Evidence for a protein kinase cascade in higher plants 3-hydroxy-3-methylglutaryl-CoA reductase kinase. *European Journal of Biochemistry* 209, 923-931.  
*Showed the existence in a number of plants, including potato, of a plant kinase able to inactivate mammalian HMG-CoA reductase. The kinase is inactivated and reactivated by mammalian protein phosphatases (2A or 2C) and mammalian kinase kinase respectively. This is probably the first evidence for a protein kinase cascade in higher plants.*
- Reymond P, Grunberger S, Paul K, Muller M, Farmer EE. 1995. Oligogalacturonide defense signals in plants: Large fragments interact with the plasma membrane *in vitro*. *Proceedings of the National Academy of Science USA* 92, 4145-4149.  
*Showed that, in vitro, elicitor-active oligogalacturonides enhanced the phosphorylation of a 34 kDa protein in purified plasma membranes from potato.*
- Seo S, Okamoto M, Seto H, Ishizuka K, Sano H, Ohashi Y. 1995. Tobacco MAP kinase: a possible mediator in wound signal transduction pathways. *Science* 270, 1988-1992.  
*Provides further experimental evidence that MAP kinases occur in plants.*
- Seo S, Niki T, Ohtsubo N, Sano H, Ohashi Y. 1997. MAP kinase in pathogen-or wound-stress in plants. *Plant Cell Physiology* 38, (supplement) S40 (SVIII-05).  
*Isolated a tobacco MAP kinase gene, WIPK, which transcript accumulated within 1 minute after wounding. Suggested the activated WIPK induces the wound-jasmonic acid signal transduction pathway.*
- Zhang S, Klessig DF. 1997. Salicylic acid activates a 48-kD MAP kinase in tobacco. *The Plant Cell* 9, 809-824.  
*Cloned a SIP (salicylic acid induced protein) kinase from tobacco; it's sequence suggested that it belonged to the MAP kinase family. They suggested that different members of the MAP kinase family are activated by different stresses.*

## **Polyamine biosynthesis**

- Bagga S, Rochford J, Klaene Z, Kuehn GD, Phillips GC. 1997. Putrescine aminopropyltransferase is responsible for biosynthesis of spermidine, spermine, and multiple uncommon polyamines in osmotic stress-tolerant alfalfa. *Plant Physiology* 114, 445-454
- Kumar A, Altabella T, Taylor MA, Tiburcio AF. 1997. Recent advances in polyamine research. *Trends in Plant Science* 2, 124-130.  
*Summarises the polyamine biosynthetic pathway showing its relationship with ethylene biosynthesis.*
- Parr AJ, Mellon FA, Colquhoun IJ, Davies HV. 2005. Dihydrocaffeoyl polyamines (kukoamine and allies) in potato (*Solanum tuberosum*) tubers detected during

metabolite profiling. *Journal of Agricultural and Food Chemistry* 53, 5461-5466.

- Ravanel S, Gakiere B, Job D, Douce R. 1998. The specific features of methionine biosynthesis and metabolism in plants. *PNAS* 95, 7805-7812.
- Stroinski A. 1997. Effects of cadmium on the host-pathogen system. V. Effect of exogenous dicyclohexylamine on potato tubers (*Solanum tuberosum* L.), cadmium and *Phytophthora infestans* relations. *Journal of Plant Physiology* 150, 178-183. *Suggested that polyamines can participate in signal transduction affecting resistance to P. infestans.*
- Torrigiani P, Rabiti AL, Bortolotti C, Betti L, Marani F, Canova A, Bagni N. 1997. Polyamine synthesis and accumulation in the hypersensitive response to TMV in *Nicotiana tabacum*. *New Phytologist* 135, 467-473. *Free and conjugated polyamines increased in resistant tobacco plants in response to infection by TMV.*
- Walden R, Cordeiro A, Tiburcio AF. 1997. Polyamines: small molecules triggering pathways in plant growth and development. *Plant Physiology* 113, 1009-1013. *Describes the biosynthesis of polyamines and shows how it is linked to ethylene biosynthesis.*
- Walters DR. 2000. Polyamines in plant-microbe interactions. *Physiological and Molecular Plant Pathology* 57, 137-146.

## **PR Proteins**

- Bol JF, Linthorst HJM, Cornelissen BJC. 1990. Plant pathogenesis-related proteins induced by virus infection. *Annual Review of Phytopathology* 28,113-138. *Provides a general summary of PR proteins with detailed information on tobacco PR proteins.*
- Boyle B, Brisson N. 2001. Repression of the defense gene PR-10a by the single-stranded DNA binding protein SEBF. *The Plant Cell* 13, 2525-2537. *SEBF = silencing element binding factor*
- Choi DS, Hwang IS, Hwang BK. 2012. Requirements of the cytosolic interaction between PATHOGENESIS-RELATED PROTEIN1) and LEUCINE-RICH REPEAT PROTEIN1) for cell death and defense signalling in pepper. *Plant Cell*. April 6.
- Desveaux D, Despres C, Joyeux A, Subramaniam R, Brisson N. 2000. PBF-2 is a novel single-stranded DNA binding factor implicated in PR-10a gene activation in potato. *Plant Cell* 12, 1477-90.
- González-Lamothe R, Boyle P, Dulude A, Roy V, Lezin-Doumbou C, Kaur GS, Bouarab K, Després C, Brisson N. 2008. *Plant Cell* 20, 3136-47.
- Kombrink E, Schroder M, Hahlbrock K. 1988. Several "pathogenesis-related" proteins in potato are 1,3-*B*-glucanases and chitinases. *Proceedings of the National*



Academy of Science USA. 85, 782-786.

- Kwon T-H, Kim T-G, Kim D-H, Yang M-S. 1999. Molecular cloning of a genomic DNA encoding the potato *B-1,3*-glucanase. *Journal of Plant Physiology* 155, 520-526.
- Shi X, Tian Z, Liu J, van der Vossen EA, Xie C. 2011. A potato pathogenesis-related protein gene, *StPRp27*, contributes to race non-specific resistance against *Phytophthora infestans*. *Mol Biol Rep* June 11.
- van Loon LC, van Strien EA. 1999. The families of pathogenesis-related proteins, their activities, and comparative analysis of PR-1 type proteins. *Physiological and Molecular Plant Pathology* 55, 85-97.  
*Describes the families of PR proteins from PR-1 to PR-14. Not specific to potatoes.*

## **Prenylation**

- Gelb MH. 1997. Protein prenylation, et cetera: signal transduction in two dimensions. *Science* 275, 1750-1751.  
*Reviewed many aspects of prenylation in eukaryotes and suggested that most prenylated proteins are members of signal transduction cascades.*
- Ohnuma S, Narita K, Nakazawa T, Ishida C, Takeuchi Y, Ohto C, Nishino T. 1996. A role of the amino acid residue located on the fifth position before the first aspartate-rich motif of farnesyl diphosphate synthase on determination of the final product. *The Journal of Biological Chemistry* 271, 30748-30754.  
*Summarises the consecutive condensation of isopentenyl diphosphates by prenyltransferase.*
- Parmryd I, Shipton CA, Andersson B, Dallner G. 1997. Protein prenylation in spinach - tissue specificity and greening-induced changes. *Archives of Biochemistry and Biophysics* 339, 73-78.  
*Provides additional experimental evidence on the importance of prenylation in post-translational modification of proteins.*
- Villar KD, Mitsuzawa H, Yang W, Sattler I, Tamanoi F. 1997. Amino acid substitutions that convert the protein substrate specificity of farnesyltransferase to that of geranylgeranyltransferase type I. *The Journal of Biological Chemistry* 272, 680-687.  
*The substrate specificity of yeast FTase can be switched to that of the closely related enzyme geranylgeranyltransferase type I (GGTase I) by a single amino acid change.*
- Quian D, Zhou D, Ju R, Cramer CL, Yang Z. 1996. Protein farnesyltransferase in plants: Molecular characterization and involvement in cell cycle control. *The Plant Cell* 8, 2381-2394.  
*Emphasises the role of isoprenylation of proteins as one factor involved in post-translational modification of proteins and subsequent control of enzyme activity. Reviewed the role of farnesyltransferase (FTase) and evidence for its role in the control of the cell cycle in plants.*

## Sesquiterpenes and phytoalexins

- Afzal M, Al-Oriquat G. 1986.  $^{13}\text{C}$  and proton-NMR spectra of sesquiterpenoid and related phytoalexins. *Heterocycles* 24, 2943-2961.  
*Provides structural data for some of the potato sesquiterpene phytoalexins.*
- Alves LM, Kirchner RM, Lodato DT, Nee PB, Zappia JM, Chichester ML, Stuart JD, Kalan EB, Kissinger JC. 1984. Acetyldehydrorishitinol, a rishitinol-related potato stress metabolite. *Phytochemistry* 23, 537-538.  
*Provides structural information on a rishitinol-related sesquiterpene induced by ethylene.*
- Bulow N, Konig WA. 2000. The role of germacrene D as a precursor in sesquiterpene biosynthesis: investigations of acid catalyzed, photochemically and thermally induced rearrangements. *Phytochemistry* 55, 141-168.
- Desjardins AE, Gardner HW, Plattner RD. 1989. Detoxification of the potato phytoalexin lubimin by *Gibberella pulicaris*. *Phytochemistry* 28,431-437.  
*Proposes a possible pathway for the degradation of lubimin by G. pulicaris involving 15-dihydrolubimin, 2-dehydrolubimin, isolubimin, cyclodehydroisolubimin, cyclolubimin, 11,12,-epoxycyclodehydroisolubimin and 11,12-epoxycyclolubimin.*
- Desjardins AE, Gardner HW, Weltring K-M. 1992. Detoxification of sesquiterpene phytoalexins by *Gibberella pulicaris* (*Fusarium sambucinum*) and its importance for virulence on potato tubers. *Journal of Industrial Microbiology* 9, 201-211.  
*Detoxification of rishitin and lubimin involve epoxidation, dehydrogenation, and cyclization. Structures of the major metabolites of rishitin and lubimin are provided.*
- Desjardins AE, McCormick SP, Plaisted RL, Brodie BB. 1997. Association between solavetivone production and resistance to *Globodera rostochiensis* in potato. *Journal of Agriculture and Food Chemistry* 45, 2322-2326.  
*A gene or genes that control solavetivone accumulation are located on potato chromosome V close to the H1 locus for resistance to G. rostochiensis, but are not at the same locus. The authors suggest that selection of nematode-resistant H1 genotypes might be facilitated by selection of tubers with high solavetivone ratios.*
- Engstrom K. 1998. Sesquiterpenoid spiro compounds from potato tubers infected with *Phoma foveata* and *Fusarium* spp. *Phytochemistry* 47, 985-990.
- Engstrom K. 1998. Post-infectional metabolites from infected potato tubers. *Phytochemistry* 49, 1585-1587.
- Ewing DF, Whitehead IM, Atkinson A, Threlfall DR. 1990.  $^1\text{H}$  NMR study of the stereochemistry of lubimin and related vetispirane sesquiterpenoids. *Journal of the Chemical Society Perkin Transactions* 2, 343-348.  
*Provided information on the stereochemistry of lubimin, 15-dihydrolubimin, and 3-hydroxylubimin.*
- Gardner HW, Desjardins AE, McCormick SP, Weisleder D. 1994. Detection of the potato phytoalexin rishitin by *Gibberella pulicaris*. *Phytochemistry* 37,1001-1005.

*Showed that a rishitin-tolerant strain of G. pulicaris metabolized rishitin to 13-hydroxyrishitin and 11,12-epoxyrishitin. Also provides <sup>1</sup>H NMR and <sup>13</sup>C NMR data for rishitin, 11,12-epoxyrishitin, 5,10-11,12-diepoxyrishitin, 5,10-epoxyrishitin and 13-hydroxyrishitin.*

- Ishiguri Y, Tomiyama K, Murai A, Katsui N, Masamune T. 1978. Toxicity of rishitin, rishitin-M-1 and rishitin-M-2 to *Phytophthora infestans* and potato tissue. *Annals of the Phytopathological Society of Japan* 44,52-56.  
*Provide structures and toxicity data for two metabolites of rishitin.*
- Katsui N, Murai A, Takasugi M, Imaizumi K, Masamune T. 1968. The structure of rishitin, a new antifungal compound from diseased potato tubers. *Chemical Communications* 43-44.  
*Provides a structure for rishitin.*
- Kuc J. 1982. Phytoalexins from the Solanaceae. In, JA Bailey, JW Mansfield, eds, *Phytoalexins*. Blackie, London. 81-105.  
*An early summary of many of the sesquiterpene phytoalexins produced by potato tubers.*
- Lyon GD. 1972. Occurrence of rishitin and phytuberin in potato tubers inoculated with *Erwinia carotovora* var. *atroseptica*. *Physiological Plant Pathology* 2, 411-416.  
*Showed that sesquiterpene phytoalexins such as rishitin and phytuberin were induced in potato tubers by the soft rot bacterium Erwinia carotovora.*
- Malmberg AG, Theander O. 1980. Two phytoalexin glycosides from potato tubers infected with *Phoma*. *Phytochemistry* 19, 1739-1742.  
*Identified four coumarins and seven isoprenoids in potato tubers infected with Phoma exigua var. foveata. These included the 7-O-B-D-glucopyranoside of isofraxidin and the sesquiterpene 2-(11,12-dihydroxy-11-methylethyl)-6,10-dimethyl-spiro[4,5]dec-6-en-8-one and its 12-O-B-D-glucopyranoside.*
- Matthews D, Jones H, Gans P, Coates S, Smith LMJ. 2005. Toxic secondary metabolite production in genetically modified potatoes in response to stress. *Journal of Agricultural and Food Chemistry* 53, 7766-7776.  
*Detected cedrol, diacetylphytuberin, ledol, and rishitin.*
- Stoessl A. 1982. Biosynthesis of phytoalexins. In: Bailey JA, Mansfield JW, eds. *Phytoalexins*. Glasgow and London, UK: Blackie 133-180.  
*Proposes a possible biosynthetic route for phytuberin and phytuberol.*
- Stoessl A, Stothers JB. 1983. Biosynthesis of antifungal stress metabolites from potato: observation of hydride shifts via the "B-hop" from incorporation of [2-<sup>2</sup>H<sub>3</sub>,2-<sup>13</sup>C]acetate and <sup>13</sup>Cmr. *Canadian Journal of Chemistry* 61, 1766-1770.  
*Provides structures for lubimin, 15-dihydrolubimin, phytuberol, rishitin, hydroxylubimin.*
- Stoessl A, Stothers JB, Ward EWB. 1976. Sesquiterpenoid stress compounds of the Solanaceae. *Phytochemistry* 15, 855-872.  
*An early review of the sesquiterpene phytoalexins isolated from the Solanaceae*

*providing chemical and physical characterisation of isolated phytoalexins together with an indication of the possible biosynthetic relationships.*

- Szafranek B, Chrapkowska K, Pawinska, Szafranek J. 2005. Analysis of leaf surface sesquiterpenes in potato varieties. *Journal of Agricultural and Food Chemistry* 53, 2817-2822.
- Szafranek B, Chrapkowska K, Waligora D, Palavinskas R, Banach A, Szafranek J. 2006. Leaf surface sesquiterpene alcohols of the potato (*Solanum tuberosum*) and their influence on colorado beetle (*Leptinotarsa decemlineata* Say) feeding. *Journal of Agricultural and Food Chemistry* 54, 7729-7734.  
*Identified ledol and kunzeaol*
- Zook MN, Chappell J, Kuc JA. 1992. Characterization of elicitor-induction of sesquiterpene cyclase activity in potato tuber tissue. *Phytochemistry* 31, 3441-3445.  
*Partially characterized the sesquiterpene cyclase of potato tubers treated with arachidonic acid.*

### **Shikimate pathway**

- Bentley R. 1990. The shikimate pathway - a metabolic tree with many branches. *Critical Reviews in Biochemistry and Molecular Biology* 25, 307-384. *Provides a critical review of enzymic steps in the shikimate pathway in plants and microorganisms.*
- Diaz J, Barcelo AR, de Caceres FM, 1997. Changes in shikimate dehydrogenase and the end products of the shikimate pathway, chlorogenic acid and lignins, during the early development of seedlings of *Capsicum annuum*. *New Phytologist* 136, 183-188. *Provides a biosynthetic pathway leading from 3-deoxy-D-arabino-heptulose 7-phosphate to shikimate.*
- Forlani G. 1997. Properties of the 5-enol-pyruvyl-shikimate-3-phosphate synthase isoforms isolated from maize cultured cells. *Journal of Plant Physiology* 50, 369-375. *Showed two isoforms of 5-enol-pyruvyl-shikimate-3-phosphate (EPSP) synthase, previously purified from maize cultured cells and shown to be functionally located in the plastid, were characterized with respect to physical and functional properties. Both enzymes are inhibited by glyphosate and not subject to feed-back regulation by aromatic amino acids or shikimate pathway intermediates. Data are consistent with the hypothesis of an isoform-based mechanism of enzyme level modulation in plant aromatic metabolism.*
- King RR, Calhoun LA. 2010. A feruloyltyramine trimer isolated from potato common scab lesions. *Phytochemistry* 71, 2187-9.
- Lodhi AH, Bongaerts RJM, Verpoorte R, Coomber SA, Charlwood BV. 1996. Expression of bacterial isochorismate synthase (EC 5.4.99.6) in transgenic root cultures of *Rubia peregriana*. *Plant Cell Reports* 16, 54-57.

- Zhao J, Last RL. 1996. Coordinate regulation of the tryptophan biosynthetic pathway and indolic phytoalexin accumulation in Arabidopsis. *The Plant Cell* 8, 2235-2244. *Describes the Arabidopsis tryptophan biosynthetic pathway from chorismate.*

## Tropane alkaloids

- Drager B. 2006. Tropinone reductases, enzymes at the branch point of tropane alkaloid metabolism. *Phytochemistry* 67, 327-337.
- Hashimoto T, Yamada Y. 1994. Alkaloid biogenesis: molecular aspects. *Annual Review of Plant Physiology and Plant Molecular Biology* 45, 257-285. *Summarises the biosynthetic route of tropane alkaloids showing that they originate from ornithine and/or arginine via ornithine decarboxylase. Putrescine N-methyltransferase (PMT) is the first committed enzyme driving the pathway away from polyamine biosynthesis to alkaloid biosynthesis. PMT has not yet been purified from potato.*
- Keiner R, Drager B. 2000. Calystegine distribution in potato (*Solanum tuberosum*) tubers and plants. *Plant Science* 150, 171-179.
- Nash RJ, Rothschild M, Porter EA, Watson AA, Waigh RD, Waterman PG. 1993. Calystegines in *Solanum* and *Datura* species and the death's-head hawk-moth (*Acherontia atropus*). *Phytochemistry* 34, 1281-1283. *Identified calystegine A3 and calystegine B2 from potato tubers and also showed that calystegine B2 was present in potato leaves. These polyhydroxylated tropane alkaloids are known to be potent inhibitors of glycosidases but there is no experimental evidence that their concentration is affected by disease or that they are part of a defence response to microorganisms.*

## Others

- Ames JB, Ishima R, Tanaka T, Gordon JI, Stryer L, Ikura M. 1997. Molecular mechanics of calcium-myristoyl switches. *Nature* 389, 198-202. *Discusses the molecular mechanism by which myristoylation of proteins may be associated with calcium-associated signal transduction processes. It provides information relevant to eukaryotic cells in general but is not specific to plants per se.*
- Avrova AO, Taleb N, Rokka V-M, Heilbronn J, Campbell E, Hein I, Gilroy EM, Cardle L, Bradshaw JE, Stewart HE, Fakim YJ, Loake G, Birch PRJ. 2004. Potato oxysterol binding protein and cathepsin B are rapidly up-regulated in independent defence pathways that distinguish R gene-mediated and field resistances to *Phytophthora infestans*. *Molecular Plant Pathology* 5, 45-56.
- Baebler Š, Krečič-Stres H, Rotter A, Kogovšek P, Cankar K, Kok EJ, Gruden K, Kovač M, Žel J, Pompe-Novak M, Ravnikar M. 2009. PVY<sup>NTN</sup> elicits a diverse gene expression response in different potato genotypes in the first 12h after inoculation. *Molecular Plant Pathology* 10, 263-275.

- Bakker E, Borm T, Prins P, van der Vossen E, Uenk G, Arens M, de Boer J, van Eck H, Muskens M, Vossen J, van der Linden G, van Ham R, Klein-Lankhorst R, Visser R, Smant G, Bakker J, Govers A. 2011. A genome-wide genetic map of NB-LRR disease resistance loci in potato. *Theor Appl Genetic* May 18.
- Beyer K, Binder A, Boller T, Collinge M. 2001. Identification of potato genes induced during colonization by *Phytophthora infestans*. *Molecular Plant Pathology* 2, 125-134.  
*Contains a table of potato genes induced in the compatible interaction. One of the enzymes listed is 3-isopropylmalate dehydrogenase which is involved in the biosynthesis of leucine. (See the reference by Daschner et al., below for related enzymes).*
- Bourque S, Dutartre A, Hammoudi V, Blanc S, Dahan J, Jeandroz S, Pichereaux C, Rossignol M, Wendehenne D. 2011. Type-2 histone deacetylases as new regulators of elicitor-induced cell death in plants. *New Phytol* June 8.
- Bouwmeester K, de Sain M, Weide R, Gouget A, Klamer S, Canut H, Govers F. 2011. The lectin receptor kinase LecRK-1.9 is a novel *Phytophthora* resistance component and a potential host target for a RXLR effector. *PLoS Pathog* 7(3): e1001327
- Chanda B, Xia Y, Mandal MK, Yu K, Sekine KT, Gao QM, Selote D, Hu Y, Stromberg A, Navarre D, Kachroo A, Kachroo P. 2011. Glycerol-3-phosphate is a critical mobile inducer of systemic immunity in plants. *Nat Genet* Mar 27.
- Daschner K, Couee I, Binder S. 2001. The mitochondrial isovaleryl-Coenzyme A dehydrogenase of *Arabidopsis* oxidises intermediates of leucine and valine catabolism. *Plant Physiology* 126, 601-612.  
*Does not mention potato specifically but provides a scheme for the catabolic pathway of leucine in Arabidopsis leading to the formation of HMG-CoA. Interestingly, one of the enzymes (3-methylcrotonyl-CoA carboxylase) has been shown by others (Schenk et al., 2000, Proc. Natl. Acad. Sci, USA 97, 11655-60) to be up-regulated by methyl jasmonate in Arabidopsis.*
- Desveaux D, Subramaniam R, Despres C, Mess JN, Levesque C, Fobert PR, Dangl JL, Brisson N. 2004. A "Whirly" transcription factor is required for salicylic acid-dependent disease resistance in *Arabidopsis*. *Dev Cell* 6, 229-40.  
*Show that the potato StWhy1 protein is a transcriptional activator of genes containing the PBF2 binding PB promoter element.*
- Edwards R. 1996. S-Adenosyl-L-methionine metabolism in alfalfa cell cultures following treatment with fungal elicitors. *Phytochemistry* 43, 1163-1169.  
*Studied responses of the methyl cycle in cell cultures to yeast-derived elicitors. Showed an increase in SAM synthetase, SAH hydrolase, adenosine nucleosidase, and adenosine deaminase activity in response to elicitors. The methyl cycle is associated with a large number of methylation reactions.*
- Engelhardt S, Boevink PC, Armstrong MR, Ramos MB, Hein I, Birch PR. 2012. Relocalization of late blight resistance protein R3a to endosomal compartments is

associated with effector recognition and required for the immune response. *Plant Cell* Dec 14.

- Ewan R, Pangestuti R, Thornber S, Craig A, Carr C, O'Donnell L, Zhang C, Sadanandom A. 2011. Deubiquitinating enzymes AtUBP12 and AtUBP13 and their tobacco homologue NtUBP12 are negative regulators of plant immunity. *New Phytol* Mar 9.
- Gallou A, Declerck S, Cranenbrouck S. 2011. Transcriptional regulation of defence genes and involvement of the WRKY transcription factor in arbuscular mycorrhizal potato root colonization. *Funct Integr Genomics* Aug 3.
- Gallou A, De Jaeger N, Cranenbrouck S, Declerck S. 2010. Fast track in vitro mycorrhization of potato plantlets allow studies on gene expression dynamics. *Mycorrhiza* 20, 201-207.
- Gao G, Jin LP, Xie KY, Qu DY. 2009. The potato *StLTPa7* gene displays a complex Ca-associated pattern of expression during the early stage of potato-*Ralstonia solanacearum* interaction. *Molecular Plant Pathology* 10, 15-27.
- Gao L, Tu ZJ, Millett BP, Bradeen JM. 2013. Insights into organ-specific pathogen defense responses in plants: RNA-seq analysis of potato tuber-*Phytophthora infestans* interactions. *BMC Genomics* 14, 340-
- Gilroy EM, Taylor RM, Hein I, Boevink P, Sadanandom A, Birch PR. 2011. CMPG1-dependent cell death follows perception of diverse pathogen elicitors at the host plasma membrane and is suppressed by *Phytophthora infestans* RXLR effector AVR3a. *New Phytologist* 190, 653-66.
- Godoy AV, Zanetti ME, Segundo BS, Casalongue CA. 2001. Identification of a putative *Solanum tuberosum* transcriptional coactivator up-regulated in potato tubers by *Fusarium solani* f. sp. *eumartii* infection and wounding. *Physiologia Plantarum* 112, 217-222.
- González-Lamothe R, El Oirdi M, Brisson N, Bouarab K. 2012. The conjugated auxin indole-3-acetic acid –aspartic acid promotes plant disease development. *Plant Cell*, Feb28.  
*Results are for Arabidopsis. Not yet shown for potato.*
- Gruden K, Strukelj B, Ravnikar M, Poljsak-Prijatelj M, Mavric I, Brzin J, Pungercar J, Kregar I. 1997. Potato cysteine proteinase inhibitor gene family: molecular cloning, characterisation and immunocytochemical localisation studies. *Plant Molecular Biology* 34, 317-323.  
*Accumulation of potato cysteine protease inhibitors (PCPIs) occurred in stem vacuoles after treatment with jasmonic acid. The PCPIs belong to the Kunitz-type soybean trypsin inhibitor family, show no homology with other cysteine proteinase inhibitor superfamilies, and also occur as a multigene family. They are postulated to play a role in resistance to insects and pathogens.*

- Hauschild R, von Schaewen A. 2003. Differential regulation of glucose-6-phosphate dehydrogenase isoenzyme activities in potato. *Plant Physiology* 133, 47-62.
- Hind SR, Pulliam SE, Veronese P, Shantharaj D, Nazir A, Jacobs NS, Stratmann JW. 2011. The COP9 signalosome controls jasmonic acid synthesis and plant responses to herbivory and pathogens. *Plant Journal* 65, 480-491.
- Kim JY, Gopal R, Kim SY, Seo CH, Lee HB, Cheong H, Park Y. 2013. PG-2, a potent AMP against pathogenic microbial strains, from potato (*Solanum tuberosum* L cv. Gogu Valley) tubers not cytotoxic against human cells. *Int J Mol Sci* 14, 4349-60.
- Kaschani F, Van der Hoorn, RAL. 2011. A model of the C14-EPIC complex indicates hotspots for a protease-inhibitor arms race in the oomycete-potato interaction. *Plant Signal Behav* 6, 109-112.
- Kelley BS, Lee SJ, Damasceno CM, Chakravarthy S, Kim BD, Martin GB, Rose JK. 2010. A secreted effector protein (SNE1) from *Phytophthora infestans* is a broadly acting suppressor of programmed cell death. *Plant J* 62, 357-66.
- King SR, McLellan H, Boevink PC, Armstrong MR, Bukharova T, Sukarta O, Win J, Kamoun S, Birch PR, Banfield MJ. 2014. *Phytophthora infestans* RXLR effector PexRD2 interacts with host MAPKKK{varepsilon} to suppress plant immune signaling. *Plant Cell* Mar 14.
- Lazar A, Coll A, Dobnik D, Baebler S, Bedina-Zavec A, Zel J, Gruden K. 2014. Involvement of potato (*Solanum tuberosum* L.) MKK6 in response to potato virus Y. *PLoS One* 9(8):e104553
- Lee S-J, Rose JKC. 2010. Mediation of the transition from biotrophy to necrotrophy in hemibiotrophic plant pathogens by secreted effector proteins. *Plant Signal Behav* 5, 769-772.
- Legendre L, Derckel JP, Wriese F, Correze C, Audran JC, Haye B, Lambert B. 1997. Evidence for the existence of cAMP in Lily plant flower tissues. *Phytochemistry* 44, 769-774.  
*The authors developed a radioimmunoassay to provide further evidence that cAMP exists in plants and may therefore function as a second messenger.*
- LiY, Tian Z, Liu J, Xie C. 2009. Comparative cDNA-AFLP analysis reveals that DL- $\beta$ -amino butyric acid induces resistance through early activation of the host-defence genes in potato. *Physiologia Plantarum* 136, 19-29.
- Ma J, Hanssen M, Lundgren K, Hernández L, Delatte T, Ehlert A, Liu CM, Schlupe H, Dröge-Laser W, Moritz T, Smeekens S, Hanson J. 2011. The sucrose-regulated Arabidopsis transcription factor bZIP11 reprograms metabolism and regulates trehalose metabolism. *New Phytol* April 27.
- McAinsh MR, Brownlee C, Hetherington AM. 1997. Calcium ions as second messengers in guard cell signal transduction. *Physiologia Plantarum* 100, 16-29.



*Reviews the role of calcium in signal transduction in guard cells.*

- McLellan H, Boevink PC, Armstrong MR, Pritchard L, Gomez S, Morales J, Whisson SC, Beynon JL, Birch PR. 2013. An RxLR effector from *Phytophthora infestans* prevents re-localisation of two plant NAC transcription factors from the endoplasmic reticulum to the nucleus. *PLoS Pathog* 9(10):e1003670.
- Mizzen CA, Yang X-J, Kokubo T, Brownell JE, Bannister AJ, Owen-Hughes T, Workman J, Wang L, Berger SL, Kouzarides T, Nakatani Y, Allis CD. 1996. The TAF<sub>II</sub>250 subunit of TFIID has histone acetyltransferase activity. *Cell* 87, 1261-1270. *Human TAF 250 and its homologs in Drosophila and yeast have histone acetyltransferase (HAT) activity.*
- Mohan S, Meiyalaghan S, Latimer JM, Gatehouse ML, Monaghan KS, Vanga BR, Pitman AR, Jones EE, Conner AJ, Jacobs JM. 2013. *GSL2* over-expression confers resistance to *Pectobacterium atrosepticum* in potato. *Theor Appl Genetics* Dec 27.
- Montesano M., Hyytiainen H, Wettstein R, Palva ET. 2003. A novel potato defence-related alcohol:NADP<sup>+</sup> oxidoreductase induced in response to *Erwinia carotovora*. *Plant Molecular Biology* 52, 177-189.
- Moreno M, Segura A, Garcia-Olmedo F. 1994. Pseudothionin-St1, a potato peptide active against potato pathogens. *European Journal of Biochemistry* 223, 135-9.
- Nakashita H, Yasuda M, Nitta T, Asami T, Fujioka S, Arai Y, Sekimata K, Takatsuto S, Yamaguchi I, Yoshida S. 2003. Brassinosteroid functions in a broad range of disease resistance in tobacco and rice. *The Plant Journal* 33, 887-898. *Shows the biosynthetic pathway from campesterol to brassinolide in tobacco. Pathway likely to occur in potato but no direct evidence yet.*
- Ni X, Tian Z, Liu J, Song B, Xie C. 2010. Cloning and molecular characterization of the potato RING finger protein gene *StRFP1* and its function in broad-spectrum resistance against *Phytophthora infestans*. *Journal of Plant Physiology* 167, 488-496.
- Nikiforova V, Kempa S, Zeh M, Maimann S, Kreft O, Casazza AP, Riedel K, Tauberger E, Hoefgen R, Hesse H. 2002. Engineering of cysteine and methionine biosynthesis in potato. *Amino Acids* 22, 259-278. *Shows the biosynthesis of cysteine and homocysteine in chloroplasts.*
- Orłowska E, Fiil A, Kirk H-G, Llorente B, Cvitanich C. 2011. Differential gene induction in resistant and susceptible potato cultivars at early stages of infection by *Phytophthora infestans*. *Plant Cell Rep.*
- Pajerowska KM, Parker JE, Gebhardt C. 2005. Potato homologs of *Arabidopsis thaliana* genes functional in defense signaling - identification, genetic mapping, and molecular cloning. *Molecular Plant-Microbe Interactions* 18, 1107-1119.
- Pauly M, Scheller HV. 2000. O-acetylation of plant cell wall polysaccharides: identification and partial characterization of a rhamnogalacturonan O-acetyltransferase from potato suspension-cultured cells. *Planta* 210, 659-667.

- Pazin MJ, Kadonaga JT. 1997. What's up and down with histone deacetylation and transcription. *Cell* 89, 325-328.  
*Reviews some of the ideas on how protein acetylation could regulate transcription. None of the cited references refer directly to plants.*
- Rawlyer AJ, Braendle RA. 2001. N-Acylphosphatidylethanolamine accumulation in potato cells upon energy shortage caused by anoxia or respiratory inhibitors. *Plant Physiology* 127, 240-251.  
*Identified NAPE in potato. NAPE known to be modulated by elicitors in other systems.*
- Restrepo S, Myers KL, del Pozo O, Martin GB, Hart AL, Buell CR, Fry WE, Smart CD. 2005. Gene profiling of a compatible interaction between *Phytophthora infestans* and *Solanum tuberosum* suggests a role for carbonic anhydrase. *Molecular Plant-Microbe Interactions* 9, 913-922.
- Rumeau D, Maher EA, Kelman A, Showalter AM. 1990. Extensin and phenylalanine ammonia-lyase gene expression altered in potato tubers in response to wounding, hypoxia, and *Erwinia carotovora* infection. *Plant Physiology* 93, 1134-1139.  
*Extensin and PAL mRNA as well as cell wall hydroxyproline levels accumulated in response to wounding and subsequent aerobic incubation but not under anaerobic incubation.*
- Ruocco M, Ambrosino P, Lanzuise S, Woo SL, Lorito M, Scala F. 2011. Four potato (*Solanum tuberosum*) ABCG transporters and their expression in response to abiotic factors and *Phytophthora infestans* infection. *Journal of Plant Physiology* 168, 2225-33.
- Saunders DG, Breen S, Win J, Schornack S, Hein I, Bozkurt TO, Champouret N, Vleeshouwers VG, Birch PR, Gilroy EM, Kamoun S. 2012. Host protein BSL1 associates with *Phytophthora infestans* RXLR effector AVR2 and the *Solanum demissum* immune receptor R2 to mediate disease resistance. *Plant Cell* 24, 3420-34.
- Senthil-Kumar M, Mysore KS. 2012.  $\delta$ -ornithine aminotransferase and proline dehydrogenase genes play a role in nonhost resistance by regulating pyrroline-5-carboxylate metabolism-induced hypersensitive response. *Plant Cell Environ* Feb10.
- Stewart D, Lyon GD, Tucker EJB. 1994. A Fourier-transform infrared spectroscopic and microscopic study of the infection of potato tubers by *Erwinia carotovora* ssp. *carotovora* in aerobic and anaerobic conditions. *Journal of the Science of Food and Agriculture* 66, 145-154.  
*Showed that starch and cell wall degradation occurs extensively in tuber tissue infected by E. carotovora. Lignification was not found to be extensively associated with infection.*
- Tasseva G, Richard L, Zachowski A. 2004. Regulation of phosphatidylcholine biosynthesis under salt stress involves choline kinases in *Arabidopsis thaliana*. *FEBS Letters* 566, 115-120.  
*Provides a simplified scheme of choline and phosphatidylcholine biosynthesis pathways in plants.*

- Tian ZD, Liu J, Wang BL, Xie CH. 2006. Screening and expression analysis of *Phytophthora infestans* induced genes in potato leaves with horizontal resistance. *Plant Cell Reports* 25, 1094-1103.
- Tameling WI, Nooijen C, Ludwig N, Boter M, Slootweg E, Goverse A, Shirasu K, Joosten MH. 2010. RanGAP2 mediates nucleocytoplasmic partitioning of the NB-LRR immune receptor Rx in the Solanaceae, thereby dictating Rx function. *Plant Cell* 22, 4176-94.
- Tian Z, He Q, Wang H, Liu Y, Zhang Y, Shao F, Xie C. 2015. The potato ERF transcription factor StERF3 negatively regulates resistance to *Phytophthora infestans* and salt tolerance in potato. *Plant Cell Physiol* Feb 13.
- Van den Burg H, Takken FLW. 2010. SUMO-, MAPK- and resistance protein-signaling converge at transcription complexes that regulate plant innate immunity. *Plant Signaling & Behavior* 5, 1-5.
- von Kampen J, Wettern M, Schulz M. 1996. The ubiquitin system in plants. *Physiologia Plantarum* 97, 618-624.  
*The role of the ubiquitin system in plant/pathogen interactions is still very poorly understood but this review summarises existing knowledge. Ubiquitin serves as a marker for targeting (post-translational modification) a protein for its subsequent degradation. Possibly involved in controlling proteins which are short-lived, denatured or abnormal, as well as some cellular regulators.*
- Wang X, Boevink P, McLellan H, Armstrong M, Bukharova T, Qin Z, Birch PR. 2015. A host KH RNA binding protein is a susceptibility factor targeted by an RXLR effector to promote late blight disease. *Mol Plant* April 29.
- Wu T, Tian Z, Liu J, Yao C, Xie C. 2009. A novel ankyrin repeat-rich gene in potato, Star, involved in response to late blight. *Biochemical Genetics* 47, 439-450.
- Wu T, Tian Z, Liu J, Xie C. 2009. A novel leucine-rich repeat receptor-like kinase gene in potato, *StLRPK1*, is involved in response to diverse stresses. *Molecular Biology Reports* 36, 2365-2374.

Reference list last modified 16 May 2015.